

The overlap number of a graph

Bill Rosgen
Centre for Quantum Technologies
National University of Singapore
bill.rosgen@nus.edu.sg

Lorna Stewart
Department of Computing Science
University of Alberta
Edmonton, Alberta, Canada
stewart@cs.ualberta.ca

August 18, 2010

Abstract

An overlap representation is an assignment of sets to the vertices of a graph in such a way that two vertices are adjacent if and only if the sets assigned to them overlap. The overlap number of a graph is the minimum number of elements needed to form such a representation. We find the overlap numbers of cliques and complete bipartite graphs by relating the problem to previous research in combinatorics. The overlap numbers of paths, cycles, and caterpillars are also established. Finally, we show the **NP**-completeness of the problems of extending an overlap representation and finding a minimum overlap representation with limited containment.

1 Introduction

All graphs we consider are finite and simple. The subgraph of $G = (V, E)$ induced by $V' \subseteq V$ is denoted by $G[V']$. For graph $G = (V, E)$ and vertex $v \in V$, $N(v) = \{u \mid (u, v) \in E\}$ is called the *open neighbourhood* of vertex v ; $N[v] = N(v) \cup \{v\}$ is the *closed neighbourhood* of vertex v . The minimum degree of any vertex in a graph G is denoted $\delta(G)$. A set of vertices is a *clique* if any two vertices in the set are adjacent. A clique is *maximal* if it is not contained in a larger clique. An *independent set* is a set of pairwise nonadjacent vertices. We denote by K_n the complete graph on n vertices. We use the notation P_n , for the path on n vertices, and C_n for the cycle on n vertices. We also need to introduce two covers of a graph: a *clique cover* is a covering of the vertices of a graph by cliques, and an *edge-clique cover* is a covering of the edges of a graph by cliques.

Two sets *overlap* if they intersect and neither set contains the other. An *overlap representation* (respectively *intersection representation*) for a graph G is an assignment of sets to the vertices of G such that two vertices are adjacent if and only if the sets assigned to them overlap (respectively intersect). The size of such a representation is the cardinality of the union of the assigned sets, and the minimum size of a representation is termed the *overlap number* (respectively *intersection number*) of the graph. The overlap number of graph G is denoted $\varphi(G)$. Every graph has an overlap representation. This follows from the fact that all graphs have intersection representations [16] (for an English translation, see [2]), and the observation that we can take an intersection representation for a graph and add a new element to each set, which causes sets to overlap if and only if they intersect. While intersection representations of graphs have been widely studied, overlap representations have received considerably less attention, even though overlapping is a natural relation of pairs of sets.

The intersection number parameter was introduced and bounded by Erdős, Goodman, and Pósa [4]. Specifically, they give a tight upper bound of $\lfloor n^2/4 \rfloor$ on the intersection number of an n -vertex graph. The **NP**-completeness of computing the intersection number is shown by Kou, Stockmeyer, and Wong [9]. In addition to these results, Raychaudhuri gives polynomial time algorithms for the intersection numbers of chordal graphs [12] and W_4 -free comparability graphs [13], where W_4 is a cycle on four vertices with a universal vertex.

For the overlap number, only a few algorithms and bounds are known. Using the simple technique of adding a new vertex to each set in the representation, the intersection number bound of [4] shows that $\varphi(G) \leq \lfloor n^2/4 \rfloor + n$ for any graph G with n vertices. It is known that any cocomparability graph G on n vertices has $\varphi(G) \leq n + 1$, since a containment representation of size $n + 1$ exists for \overline{G} in which every set has a common element [6], and such a representation is an overlap representation for G . Thus $K_{\frac{n}{2}, \frac{n}{2}}$ has intersection number $\lfloor n^2/4 \rfloor$ [4] and overlap number at most $n + 1$ since it is a cocomparability graph. Any graph with at most a linear number of maximal cliques must have linear intersection number, as a minimum intersection representation and a minimum edge-clique cover have the same size [4]. This implies that a graph with a linear number of maximal cliques must also have linear overlap number, since we can add a new element to each set of a minimum intersection representation. This technique yields linear bounds on the overlap number of trees, chordal graphs, and planar graphs, using previous linear bounds on the number of maximal cliques in chordal graphs [5] and planar graphs [11]. Henderson [7] gives lower bounds on the overlap number of a graph in terms of its independent sets, and constant factor approximation algorithms for the overlap numbers of trees and planar graphs. In addition, he shows that there exist graphs with overlap number quadratic in the number of vertices of the graph. Cranston et al [3] show how to compute the overlap number of a tree in linear time, and give upper bounds on the overlap numbers of some graphs. Their results include the following bounds which are satisfied with equality for some graphs: $\varphi(G) \leq |E| - 1$ where $G = (V, E)$, $G \neq K_3$, and $\delta(G) \geq 2$, and $\varphi(G) \leq n^2/4 - n/2 - 1$ where G is an n -vertex graph with $n \geq 14$.

In this paper we present hardness results for problems related to finding the overlap number, and give formulas and describe algorithms for the overlap numbers of some simple graphs. These results appeared in the first author's Master's thesis [14].

The remainder of this paper is organized as follows. In Section 2 we formally introduce the overlap number and discuss some of the basic properties of overlap representations. We follow this with Section 3, where we give formulas and describe algorithms for the overlap numbers of some graphs. Finally, in Section 4, we present some **NP**-completeness results on problems related to the overlap number.

2 The Overlap Number of a Graph

Before formalizing the definition of an overlap representation, we note that by a *collection* we refer to a multiset of sets, and for simplicity we allow the mapping from sets of a representation to vertices of a graph to remain implicit, with the set S_v associated with the vertex v of the graph. With these notational issues resolved, we define an overlap representation in the following way.

Definition 1. Given a graph $G = (V, E)$, a collection $\mathcal{C} = \{S_v : v \in V\}$ is an *overlap representation* for G if for every $u, v \in V$ we have

$$(u, v) \in E \text{ if and only if } S_u \cap S_v \neq \emptyset, S_u \not\subseteq S_v, \text{ and } S_v \not\subseteq S_u.$$




Figure 1: Example of a minimum overlap representation.

We define the *size* of a representation to be the number of elements used in the representation, which is $|\bigcup_{v \in V} S_v|$, and we let the *overlap number*, $\varphi(G)$, be the size of a minimum overlap representation.

As can be seen from the definition, overlap representations have many similarities to intersection representations: sets assigned to adjacent vertices must intersect and disjoint sets map to nonadjacent vertices. In the overlap case, however, the situation is more complex, as not only do we need to ensure a stronger condition than intersection for adjacent vertices, we have a choice of representation, for every non-edge, of disjointedness or containment. As an example, consider the representation in Figure 1, where there are nonadjacent vertices represented in each of these ways.

In the case of an intersection representation, if we take an element of the representation and examine all those sets it is contained in, we find that the vertices associated with them form a clique. Doing the same in an overlap representation once again leaves us with a collection of vertices with intersecting sets, except here we may have non-edges represented by containment, and so, since the orientation implied by set containment forms a partial order, we can map elements of the representation to cocomparability graphs. Unfortunately, while covering all edges of a graph with cliques leads to an intersection representation, if we simply cover the edges of a graph with cocomparability graphs, we do not generally end up with a valid overlap representation (most cocomparability graphs do not have overlap number one).

Observation 2. *If $\{S_v : v \in V\}$ is an overlap representation for $G = (V, E)$ then, for any $V' \subseteq V$, $\{S_v : v \in V'\}$ is an overlap representation for $G[V']$. Thus $\varphi(G) \geq \varphi(H)$ for all induced subgraphs H of G .*

Vertex multiplication is the expansion of a vertex into an independent set, such that the vertices of the independent set have the same adjacencies as the original vertex.

Observation 3. *If there is an overlap representation of size s for graph G then there is an overlap representation of size s for graph G' , where G' arises from G by vertex multiplication.*

This can be observed by simply duplicating the set assigned to a vertex when it is multiplied. Note that the size of an intersection representation is preserved by the operation of expanding a vertex to a clique but not by vertex multiplication.

We conclude this section with three results that will be used in the next section.

Lemma 4. *If A , B , and C are three sets such that $A \subseteq C$, A overlaps B , but B does not overlap C , then $B \subseteq C$.*

Proof. Since A and B overlap, we have $A \cap B \neq \emptyset$ and $A \setminus B \neq \emptyset$ which, combined with the fact that $A \subseteq C$, imply $C \cap B \neq \emptyset$ and $C \setminus B \neq \emptyset$. Now since B does not overlap C , $B \subseteq C$. \square

We amplify this lemma to the following stronger result that we use in Section 3.3 to argue bounds on the size of an overlap representation for a graph, based on the size of representations for the connected components of the graph.

Lemma 5. *Let $G = (V, E)$ be a graph with overlap representation $\mathcal{C} = \{S_v : v \in V\}$. Let X and Y be nonempty subsets of V such that each of $G[X]$ and $G[Y]$ is connected, and no edge of E has one endpoint in X and the other in Y . Let $U_X = \bigcup_{x \in X} S_x$ and $U_Y = \bigcup_{y \in Y} S_y$. If $S_x \subseteq S_y$ for some $x \in X$, $y \in Y$, then*

- (i) *for all $y \in Y$, $U_X \subseteq S_y$ or $U_X \cap S_y = \emptyset$, and*
- (ii) *if $|X| > 1$ or $|Y| > 1$ then for all $x \in X$, $y \in Y$, $S_y \not\subseteq S_x$.*

Proof. We first show that, for all $x \in X, y \in Y$,

- (1) if $S_x \subseteq S_y$ then $U_X \subseteq S_y$, (2) if $S_y \subseteq S_x$ then $U_Y \subseteq S_x$, and (3) if $S_x = S_y$ then $|X| = |Y| = 1$.

Suppose (1) is false. Let $x' \in X$ be such that $S_{x'}$ contains an element that is not in S_y , where the distance in $G[X]$ from x to x' is as small as possible. Let $x = x_1 x_2 \dots x_k = x'$ be a shortest x, x' -path in $G[X]$. Then $S_{x_{k-1}} \subseteq S_y$ (by the choice of x'), $S_{x_{k-1}}$ overlaps $S_{x'}$ (because x_{k-1} and x' are adjacent on the path), and $S_{x'}$ does not overlap S_y (since x' and y are not adjacent in G). But then by Lemma 4, $S_{x'} \subseteq S_y$, contradicting the choice of x' . The justification for (2) is similar. For (3), if $S_x = S_y$ then x and y are adjacent to exactly the same vertices in G , and therefore, since each of $G[X]$ and $G[Y]$ is connected, and there are no edges between X and Y in G , $|X| = |Y| = 1$.

Note that (i) is true if $|X| = |Y| = 1$ since then $U_X = S_x \subseteq S_y$ for $x \in X, y \in Y$. Now suppose (i) is false. Then $|X| > 1$ or $|Y| > 1$ and there exists $y \in Y$ with $U_X \not\subseteq S_y$ and $U_X \cap S_y \neq \emptyset$. By (1), there is no $x \in X$ with $S_x \subseteq S_y$. Therefore, $S_y \subseteq S_{x'}$ for some $x' \in X$. We also have $S_x \subseteq S_{y'}$ for some $x \in X, y' \in Y$, by the statement of the lemma. But now, by (1) and (2), $U_X \subseteq S_{y'} \subseteq U_Y \subseteq S_{x'} \subseteq U_X$, which implies $S_{x'} = S_{y'}$, contradicting (3).

If (ii) is false, we again have $S_x \subseteq S_{y'}$ and $S_y \subseteq S_{x'}$ for some $x, x' \in X$ and $y, y' \in Y$ which, together with (1) and (2), contradicts (3). \square

In Section 3.2 we use the following simplification of Lemma 5 to argue bounds on the overlap numbers of paths, cycles, and caterpillars.

Corollary 6. *Let $G = (V, E)$ be a graph, and let $\mathcal{C} = \{S_v : v \in V\}$ be an overlap representation of G . Fix $v \in V$, and let, for $u \in V \setminus N[v]$, $A_v(u)$ be the vertex set of the connected component of $G[V \setminus N[v]]$ that contains u . If $S_u \subseteq S_v$, then $\bigcup_{w \in A_v(u)} S_w \subseteq S_v$.*

3 Minimum Overlap Representations

In this section, we give formulas for the overlap numbers of cliques, complete k -partite graphs, paths, cycles, and caterpillars and for the overlap number of a disconnected graph in terms of the overlap numbers of its connected components.

3.1 Cliques and Complete k -partite Graphs

An overlap representation of a clique is simply a collection of sets where no set contains any other and no two sets are disjoint. We can apply a theorem of Milner to find the minimum size of such a representation.

Definition 7. The maximum size of a family, \mathcal{C} , of subsets of $\{1, 2, \dots, m\}$ satisfying, for $p \geq 0$,

1. If $A, B \in \mathcal{C}$, with $A \neq B$, then $A \not\subseteq B$,
2. If $A, B \in \mathcal{C}$, then $|A \cap B| \geq p$,

is denoted $S(p, m)$.

The value of the function $S(p, m)$ is exactly the quantity given by Milner's Theorem, first published in 1966.

Theorem 8 (Milner [10]). *For $m \geq 1$ and $p \geq 0$,*

$$S(p, m) = \binom{m}{\lfloor \frac{m+p+1}{2} \rfloor}.$$

Also noted by Milner [10], is that it is easy to construct a collection that achieves this bound, by simply choosing all subsets of $\{1, 2, \dots, m\}$ of size $\lfloor (m+p+1)/2 \rfloor$. This, reformulated in the language of overlap representations, is precisely the content of the following corollary.

Corollary 9. *For $n \geq 1$, $\varphi(K_n) = \min \{m : n \leq S(1, m)\}$.*

Proof. Consider any overlap representation, \mathcal{B} , of K_n . Any two elements of \mathcal{B} must intersect, and no element can contain any other, as each pair of vertices in K_n forms an edge. Thus $|\mathcal{B}| \leq S(1, m)$, where $m = |\bigcup_{A \in \mathcal{B}} A|$.

For any m , consider the collection given by $\mathcal{C}_m = \{A \subseteq \{1, 2, \dots, m\} : |A| = \lfloor (m+2)/2 \rfloor\}$. As we have $\lfloor (m+2)/2 \rfloor \geq \lceil m/2 \rceil$, any two elements of \mathcal{C}_m form an intersecting pair, and furthermore, no element is contained in any other, as they all have the same size. Counting the number of ways to form subsets of $\{1, 2, \dots, m\}$, we obtain

$$|\mathcal{C}_m| = \binom{m}{\lfloor \frac{m+2}{2} \rfloor} = S(1, m).$$

Then, to find the minimum representation, we seek the minimum m that leaves enough room to form an overlap representation. We can simply choose any n elements of \mathcal{C}_m to obtain an overlap representation on m elements. Thus, $\varphi(K_n)$ is the smallest m such that $n \leq |\mathcal{C}_m| = S(1, m)$, as desired. \square

The next result follows immediately from Corollary 9 and Observation 3.

Corollary 10. *If G is a complete k -partite graph, then $\varphi(G) = \min\{m : k \leq S(1, m)\}$.*

We now investigate some computational issues involved in finding overlap representations of cliques and k -partite graphs. This is done by first finding bounds on $\varphi(K_n)$ in terms of n which, together with the constructive proof of Corollary 9, yield a simple polynomial time algorithm to produce a minimum overlap representation of K_n .

In order to gain a view of the size of the required representation for a given graph, we unwind the expression

$$\min \left\{ m : n \leq \binom{m}{\lfloor \frac{m+2}{2} \rfloor} \right\},$$

to obtain an asymptotically tight bound on $\varphi(K_n)$ in terms of n . We make use of Stirling's Approximation, which can be found, for example, in [1]:

$$\sqrt{2\pi n}(n/e)^n \leq n! \leq e^{1/(12n)}\sqrt{2\pi n}(n/e)^n. \quad (1)$$

This results in the following lemma.

Lemma 11. *For $1 \leq k < n$,*

$$\binom{n}{k} \geq \sqrt{\frac{1}{8\pi k}} \left(\frac{n}{k}\right)^k \left(\frac{n}{n-k}\right)^{n-k}.$$

Proof. The inequality follows by substituting equation (1) into the expansion of the binomial coefficient. \square

Using this lemma, we bound the size of the minimum overlap representation of the graphs we have considered. The proof here is simply a calculation and is omitted.

Theorem 12. *For $n \geq 1$,*

$$\min \left\{ m : n \leq \binom{m}{\lfloor \frac{m+2}{2} \rfloor} \right\} \in \Theta(\log n).$$

The next result is the final ingredient needed to build an efficient algorithm to find, for a given n , the minimum m such that $n \leq S(1, m)$.

Proposition 13. *For any $m \geq 2$,*

$$S(1, m) = \begin{cases} \frac{2m}{m-1} S(1, m-1) & \text{if } m \text{ is odd,} \\ \frac{2m}{m+2} S(1, m-1) & \text{if } m \text{ is even.} \end{cases}$$

Proof. The proof makes use of the following identities on binomial coefficients

$$\begin{aligned} \binom{n}{k} &= \frac{n}{n-k} \binom{n-1}{k} \\ \binom{n}{k} &= \frac{n}{k} \binom{n-1}{k-1}, \end{aligned}$$

which can be found, for example, in [8]. \square

Using the recurrence of Proposition 13, we can compute $S(1, m)$ for successive values of m , until we find $\varphi(K_n)$, the smallest m such that $n \leq S(1, m)$. By Theorem 12, this produces an algorithm with runtime in $O(\log n)$. To compute a minimum representation for K_n , as noted in the proof of Corollary 9, we simply take any n of the subsets of $\{1, 2, \dots, \varphi(K_n)\}$ of cardinality $\lfloor (\varphi(K_n) + 2)/2 \rfloor$. Since $\varphi(K_n) \in O(\log n)$ (by Theorem 8, Corollary 9 and Theorem 12), n of these subsets can be found in $O(n)$ time. These algorithms can also be immediately extended to find representations for complete n -partite graphs, as described in Observation 3.

3.2 Paths, Cycles, and Caterpillars

A minimum intersection representation for a path is simple to find, as there is only one possible edge-clique cover, the one consisting of each maximal clique. While it is essentially no harder to find an overlap representation of a path, proving the optimality of the representation is more difficult. Once we have shown the size that an overlap representation of a path must have, we extend the result to the case of cycles and caterpillars. The construction, given as part of the proof of the following theorem can be immediately transformed into a efficient algorithm for generating an overlap representation of P_n . This theorem does not hold for P_2 , as $\varphi(P_2) = 3$.

Theorem 14. *For $n \geq 3$, $\varphi(P_n) = n$.*

Proof. For $n = 3$, we observe that $\{\{1, 2\}, \{2, 3\}, \{1, 2\}\}$ is a minimum overlap representation, since we need at least three elements to represent a single edge. We now show that, for $n \geq 4$, $\varphi(P_n) \geq \varphi(P_{n-1}) + 1$, thereby proving that $\varphi(P_n) \geq n$. For $n \geq 4$, let $1, 2, \dots, n$ be the vertices of P_n in the order in which they appear on the path, and let $\mathcal{C} = \{S_1, S_2, \dots, S_n\}$ be a minimum overlap representation for P_n . By Corollary 6, either S_1 contains none of $\{S_3, S_4, \dots, S_n\}$, or it contains each S_i for $i \geq 3$. Notice also that these two cases collapse, since if S_1 contains all S_i for $i \geq 3$, then in particular, $S_n \subseteq S_1$, and so, if we consider the reversal of the path, we find that S_n contains none of the other sets, since $n \geq 4$. We thus need only consider the first case.

To this end, let the representation, without loss of generality, be such that the set S_1 contains none of $\{S_3, \dots, S_n\}$. Notice that with the exception of S_2 , the elements of S_1 are either all contained in one of the other sets, or none of them are. We form a representation for P_{n-1} where these elements are compressed into a single element. We consider the collection given by $\mathcal{C}' = \{S_2 \cup S_1, S_3, \dots, S_n\}$. As S_1 and S_2 share at least one element, S_k does not overlap $S_2 \cup S_1$ for any $k \geq 4$. To see this, we consider two cases. The first case is that $S_1 \subseteq S_k$, but then, by Lemma 4, we have $S_2 \subseteq S_k$ as well, which implies that $S_1 \cup S_2 \subseteq S_k$, as desired. In the other case we have S_1 disjoint from S_k , but in this case we can observe that $S_2 \not\subseteq S_k$, as this would imply, again by Lemma 4, that $S_1 \subseteq S_k$. Since $S_2 \not\subseteq S_k$, it is either disjoint from S_k , in which case $S_1 \cup S_2$ is as well, or $S_k \subseteq S_2$, which implies that $S_k \subseteq S_1 \cup S_2$, as required. Similarly, in the collection $\mathcal{C}'' = \{S_2 \setminus S_1, S_3, \dots, S_n\}$, a case analysis shows that the set $S_2 \setminus S_1$ does not overlap any set S_k for $k \geq 4$.

Thus, we need only verify that one of these two collections preserves the overlap between S_3 and the replacement for S_2 . To see that at least one suffices, let \mathcal{C}' fail to be an overlap representation for P_{n-1} , which implies that $S_3 \subseteq S_2 \cup S_1$, as we have only enlarged S_2 . Thus, $S_1 \cap S_3 \neq \emptyset$, as S_3 is contained in neither S_1 or S_2 , but it is contained in their union. Also, since S_1 does not contain any other set in the representation, we must have $S_1 \subseteq S_3$ as these two vertices are not adjacent in the path. Notice also that, since S_3 is contained in $S_1 \cup S_2$, and S_3 is not contained in S_1 , we must have $S_3 \cap (S_2 \setminus S_1) \neq \emptyset$. Seeking a contradiction, we assume that S_3 and $S_2 \setminus S_1$ also do not overlap. Since \mathcal{C} is an overlap representation for P_n , we must have $S_3 \not\subseteq S_2 \setminus S_1 \subseteq S_2$,

as the vertices associated with S_2 and S_3 are adjacent. This leaves only one way for S_3 to fail to overlap $S_2 \setminus S_1$, which is $(S_2 \setminus S_1) \subseteq S_3$. If this is the case, then we have $S_2 \subseteq S_3$, as we know that $S_1 \subseteq S_3$, which we derived from the failure of \mathcal{C}' . This contradicts the fact that \mathcal{C} is an overlap representation for P_n , and so one of \mathcal{C}' and \mathcal{C}'' must form a valid representation for P_{n-1} . In both of these representations, each set either contains S_1 or is disjoint from it, and so there is no loss in replacing the elements of S_1 with a single element. This reduces the size of the representation by at least one, as a set needs at least two elements to overlap another set. Hence, we have formed a representation of P_{n-1} of size at most $\varphi(P_n) - 1$. By induction on n , we have shown that

$$\varphi(P_n) \geq 1 + \varphi(P_{n-1}) = 1 + n - 1 = n.$$

To finish the proof, it is sufficient to build a representation of this size. Consider the representation for P_n given by, for $1 \leq i \leq n - 1$,

$$\begin{aligned} S_i &= \{i, i + 1\} \\ S_n &= \{1, 2, \dots, n - 1\}. \end{aligned}$$

Notice that in this representation, on the first $n - 1$ vertices, the set S_i overlaps only the sets S_{i-1} and S_{i+1} and is disjoint from the other sets, with the exception of S_n . Also, S_n contains all sets except S_{n-1} , which it overlaps, and so this is an overlap representation for P_n using n elements. This proves that $\varphi(P_n) = n$. \square

The representation used in the proof of the theorem is optimal in the number of elements used, and can be constructed in $O(n)$ time, which is asymptotically optimal, as a representation needs to have linear size. Thus we can view this construction as an efficient algorithm to find an overlap representation of a path.

Having found the overlap number of a path, we can find immediate lower bounds on the size of the overlap representation for some other simple graphs. The first of these is C_n , the cycle on n vertices. Once again, the lower bound is matched by a simple construction, which can be transformed immediately into an algorithm with running time linear in n . This result is not true for $n = 3$, as $\varphi(C_3) = 3$.

Corollary 15. *For $n \geq 4$, $\varphi(C_n) = n - 1$.*

Proof. To see that $\varphi(C_n) \geq n - 1$ we simply observe that by Theorem 14, the size of the representation for any $n - 1$ of the n vertices is at least $n - 1$, and so it remains only to construct a representation using $n - 1$ elements. We do this by setting, for $1 \leq i \leq n - 2$, $S_i = \{i, i + 1\}$, which forms an overlap representation for a path of $n - 2$ vertices, using $n - 1$ elements. We add to this representation $S_{n-1} = \{1, 2, 3, \dots, n - 2\}$ and $S_n = \{2, 3, 4, \dots, n - 1\}$, noting that S_{n-1} overlaps only S_n and S_{n-2} , containing the other sets, and that S_n overlaps only S_{n-1} and S_1 as it contains all other sets in the collection. Thus, the collection $\mathcal{C} = \{S_1, S_2, \dots, S_n\}$ forms an overlap representation for C_n using $n - 1$ elements, proving that $\varphi(C_n) = n - 1$. \square

We next consider overlap representations of caterpillars. A tree is a *caterpillar* if the non-leaf vertices form a path, known as the *spine* of the caterpillar. We use Theorem 14 to find a lower bound on the size of an overlap representation for a caterpillar, and pair this result with a simple construction to show that the bound is tight.

Corollary 16. *For a caterpillar T with spine containing $k \geq 1$ vertices, $\varphi(T) = k + 2$.*

Proof. We show that the size of a minimum overlap representation for a caterpillar has size determined by the size of the overlap representation for the longest path in the caterpillar. Let T be a caterpillar, and label the vertices of the spine in order $\{1, 2, \dots, k\}$, and let L_i be the leaves connected to vertex i of the spine. Notice that any longest path in T has a vertex in L_1 and a vertex in L_k as endpoints, with the remaining vertices being those of the spine. This allows the above labelling scheme to be implemented in linear time, as a longest path in a tree can be found in linear time. Also notice that the longest path in T contains $k + 2$ vertices, and so Theorem 14 provides a lower bound of $\varphi(T) \geq k + 2$.

To show a tight bound, we need only find a representation of the correct size. The representation used is similar to the one used in the proof of Theorem 14. For T a caterpillar, with nodes labelled $1, 2, \dots, k$ that form the spine, with node i adjacent to nodes $i - 1$ and $i + 1$, and L_i the set of leaves adjacent to vertex i , consider the representation given by, for $1 \leq i \leq k$,

$$\begin{aligned} S_i &= \{i + 1, i + 2\} \\ S_{L_i} &= \{1, 2, \dots, i + 1\}, \end{aligned}$$

where the set S_{L_i} is associated with all vertices in L_i . This representation coincides with the one previously given for paths, since viewing a path on n vertices as a caterpillar produces a caterpillar with $n - 2$ vertices on the spine, and two leaves, one on each end of the path. To see that the given representation is correct, notice that two vertices of the spine i and j overlap if and only if $|i - j| = 1$. Notice also that the sets assigned to two leaves never overlap, as S_{L_i} overlaps all S_{L_j} for $j \leq i$. In addition, S_{L_i} overlaps only S_i , since S_{L_i} contains S_j for $j < i$, and S_{L_i} is disjoint from S_j for $j > i$. This proves that $\varphi(T) = k + 2$. \square

This representation can be efficiently constructed in the sum of the sizes of the sets of the representation, which is $O(nk)$.

3.3 Disconnected Graphs

In this section we examine the size of a minimum overlap representation for a disconnected graph based on the sizes of minimum overlap representation of each of the connected components of the graph. This allows us to find the minimum overlap representation of a graph composed of the pieces we have already studied, and may lead to divide and conquer algorithms to find the size of an overlap representation for graphs such as threshold graphs and cographs that can be defined in terms of decomposition schemes.

Theorem 17. *If G is a graph with connected components B_1, B_2, \dots, B_k , then*

$$\varphi(G) = \sum_{i=1}^k \varphi(B_i) - (k - 1).$$

Proof. If $k = 1$, the theorem is trivially true. We assume that all components of G have size at least two, as isolated vertices can be added to a nonempty graph without increasing the size of the overlap representation, by assigning the isolated vertex a set consisting of any single element. In the case that G consists only of isolated vertices, the theorem is also trivially true. To prove this

theorem we first, as before, show a lower bound, and then argue that a representation achieving this lower bound must exist.

Suppose $k = 2$. By Lemma 5, the two components must either be independent, with no elements in common in the overlap representation, or some sets of one component can contain all sets of the other. If the two components are independent then $\varphi(G) = \varphi(B_1) + \varphi(B_2)$. In the other case, assume without loss of generality that some set associated with a vertex of B_2 contains a set associated with a vertex of B_1 . Thus, by Lemma 5, any set associated with a vertex of B_2 that intersects the set U of elements in the union of the sets associated with the vertices of B_1 , must contain all of U . In this case the elements of U may be considered to act as a single element and so, given a minimum overlap representation for G , we can take the representation restricted to B_2 and replace the elements of U by a single new element, resulting in an overlap representation for B_2 of size $\varphi(G) - \varphi(B_1) + 1$. Therefore, $\varphi(B_2) \leq \varphi(G) - \varphi(B_1) + 1$, which yields the desired bound of

$$\varphi(G) \geq \varphi(B_1) + \varphi(B_2) - 1.$$

In the case that $k \geq 3$, we consider a minimum overlap representation $\mathcal{C} = \{S_v : v \in V\}$, and once again show a lower bound on the size of \mathcal{C} . Take any three components with vertex sets A, B , and C . If some set associated with a vertex of A is contained in a set, S_b for $b \in B$, and some set associated with not necessarily the same vertex of A is contained in S_c for $c \in C$, then, by Lemma 5 the sets S_b and S_c must contain $\bigcup_{a \in A} S_a$. In particular, S_b and S_c intersect, and so one set must contain the other, as they are sets associated with nonadjacent vertices in G . This forces a containment relationship between B and C , so that the set associated with any vertex of A is forced to be contained in the sets associated with the vertices of one of B or C by transitivity. To see how this observation is useful, we build a graph F' , where the vertices of the graph are components in G , and two vertices A and B are connected by a directed edge if there is some vertex $a \in A$ and $b \in B$ such that $S_a \subseteq S_b$ in \mathcal{C} . Notice that by Lemma 5, each pair of vertices is either nonadjacent, or connected by one directed edge. The above observation is then simply the observation that no vertex, v , of F' is connected to two nonadjacent vertices by edges directed away from v . This implies that if we take the transitive reduction of F' , we obtain a graph with no cycles, and this graph remains acyclic even if we discard the orientation of the edges. Let F be the directed forest resulting from this transitive reduction. Since the edges of F represent containment and no vertex is connected by directed edges to two nonadjacent vertices, each tree has a unique root that all edges of the tree are directed towards.

As in the case that $k = 2$, if two components B_i and B_j are related by containment such that $S_i \subseteq S_j$ for some $i \in B_i, j \in B_j$, the elements of $U = \bigcup_{v \in B_i} S_v$ function as a single element in the representation for the vertices of B_j , which is otherwise unrestricted. Thus if we take an overlap representation of these two components we are able to find a representation that is at most one element smaller than the representation of the two components by disjoint sets. Notice that we can save this one element once for every edge of F , as these edges count exactly the containment relationships that are not forced by transitivity. The largest number of edges F can have is one fewer than the number of components of G , as there must be some root vertex that is not connected by a directed edge to any other vertex. This provides the following lower bound,

$$\varphi(G) \geq \sum_{i=1}^k \varphi(B_i) - (k - 1). \tag{2}$$

To show that a representation exists that achieves this bound, we take a minimum overlap representation for each component B_i of G , such that any two of these representations are disjoint. We then, for each i in increasing order, create a containment relationship between B_i and B_{i+1} , by choosing an arbitrary element of the representation for B_{i+1} and replacing it with the union of all elements used in the representation of B_i . The resulting representation is a valid overlap representation for G , as we have replaced elements in such a way as to not affect the overlapping properties within a component, and, given any two components, if two sets of the representations associated with them have nonempty intersection, then one set must contain the other, so that there are no adjacencies created between components. Notice that this representation has size given by Equation (2), as we have taken optimal representations for each component, and removed exactly $k - 1$ elements, and so this is an optimal overlap representation for G , of size $\sum_{i=1}^k \varphi(B_i) - (k - 1)$, which proves the theorem. \square

4 Hardness Results

In this section we present some **NP**-completeness results for problems related to finding the minimum overlap representation of a given graph.

4.1 Extending a Representation

A natural approach to finding the overlap number for a graph is to employ a greedy strategy, adding one vertex at a time, and only making changes to the set associated with the newly added vertex. Unfortunately, this is not a feasible approach for a general graph and overlap representation, as the problem of deciding whether or not a new element needs to be added to the representation is **NP**-complete. The formal statement of this decision problem is as follows.

Problem. The OVERLAP EXTENSION problem is defined as:

Instance: A graph, $G = (V, E)$, an overlap representation $\mathcal{C} = \{S_v : v \in V\}$ of G , and a set $A \subseteq V$.

Question: Is there a set $S \subseteq \bigcup_{v \in V} S_v$ that overlaps S_v if and only if $v \in A$?

Since such an extension can be efficiently verified, this problem is in **NP**. To see that the related problem on intersection representations can be solved efficiently, notice that in the intersection case, an element i of the representation can be added to S if and only if the set A contains all vertices v with $i \in S_v$. If all elements that can be added to S fail to form an intersection representation for the extended graph, then without introducing a new element, no such extension is possible.

Returning to the overlap case, the problem that we reduce to OVERLAP EXTENSION is the NOT-ALL-EQUAL 3SAT problem, which is identical to the standard 3SAT problem, with the exception that we seek a satisfying truth assignment where no clause has all true literals. This problem is known to be **NP**-complete [15].

Theorem 18. OVERLAP EXTENSION is **NP**-complete.

Proof. Let (U, F) be an instance of NOT-ALL-EQUAL 3SAT, where $U = \{x_1, x_2, \dots, x_n\}$ is the set of variables, and $F = \{c_1, c_2, \dots, c_m\}$ is the set of clauses with $|c_i| = 3$, for each i . If $n < 4$, we can

examine all possible truth assignments to determine if there is a solution to the NOT-ALL-EQUAL 3SAT instance, and output a trivial yes or no instance of OVERLAP EXTENSION.

If $n \geq 4$, we construct a graph $G = (V, E)$, an overlap representation \mathcal{C} of G , and a set A , to form an instance of OVERLAP EXTENSION. The vertices in the graph are given by

$$V = \{v_i : 1 \leq i \leq n\} \cup \{w_i : 1 \leq i \leq m\},$$

where each v_i is associated with a variable $x_i \in U$, and each w_i is associated with a clause $c_i \in F$. We take the overlap representation representation \mathcal{C} given by

$$\mathcal{C} = \{S_{v_i} = \{x_i, \neg x_i\} : x_i \in U\} \cup \{S_{w_i} = c_i : c_i \in F\},$$

and we set E to those edges consistent this representation. Finally, we let $A = V$ to complete the instance (G, \mathcal{C}, A) of OVERLAP EXTENSION. This transformation can clearly be performed in polynomial time. A solution of the extension problem is a set of literals that overlaps each set in \mathcal{C} , and we show that such a set is equivalent to a satisfying truth assignment for (U, F) in which each clause has at least one false literal.

To see this, let $S \subseteq U \cup \{\neg x : x \in U\}$ be a set that overlaps all elements of \mathcal{C} . Since S overlaps each $S_{v_i} = \{x_i, \neg x_i\}$, S must contain exactly one element of S_{v_i} , and so we consider the truth assignment T that makes each literal in S true. In addition, S overlaps each $S_{w_i} = c_i$, which forces at least one, but not all, of the literals in c_i to be contained in S , which shows that T satisfies the clause c_i without making all literals true.

In the other direction, we take any truth assignment T that satisfies (U, F) without making all literals in any clause true, and consider the set S of all literals made true by T . Since T is a truth assignment, for each $1 \leq i \leq n$, S contains exactly one of x_i and $\neg x_i$, and so S overlaps S_{v_i} for all i . Furthermore, since T is a satisfying truth assignment, S must intersect each S_{w_i} , and it cannot contain any S_{w_i} , as this would imply that T satisfies all literals of each clause c_i . Finally, $|S_{w_i}| = |c_i| = 3$, and $|S| = |U| \geq 4$, so S_{w_i} cannot contain S for any i . This implies that S overlaps S_{w_i} for all $1 \leq i \leq m$, and so S is a solution to the instance of OVERLAP EXTENSION. \square

Using a similar reduction, we can show the hardness of the problem of the CONTAINMENT EXTENSION problem, which is the analogue of the OVERLAP EXTENSION problem on containment representations. In this case the reduction is from the well-known **NP**-complete 3SAT problem.

Theorem 19. CONTAINMENT EXTENSION is **NP**-complete.

Proof. Let (U, F) be an instance of 3SAT, where $U = \{x_1, x_2, \dots, x_n\}$ is a set of n variables, and $F = \{c_1, c_2, \dots, c_m\}$ is a set of m clauses, each containing three literals. We may once again consider only the case where $n \geq 4$, as the reduction can output a trivial yes or no instance if this is not the case.

The vertices of the constructed graph $G = (V, E)$ are given, similarly to the OVERLAP EXTENSION case, by

$$V = \{v_i : 1 \leq i \leq n\} \cup \{w_i : 1 \leq i \leq m\} \cup \{z\}.$$

We set $L = \bigcup_{i=1}^n \{x_i, \neg x_i\}$, the set of all literals, and construct the containment representation given by the collection \mathcal{C} consisting of the following sets, for all $1 \leq i \leq n$ and $1 \leq j \leq m$,

$$\begin{aligned} S_{v_i} &= \{x_i, \neg x_i\} \\ S_{w_i} &= (L \setminus c_i) \cup \{0\} \\ S_z &= \{0\}. \end{aligned}$$

To complete the constructed instance, we set $A = \{z\}$.

In a similar way to the proof of Theorem 18, it can be observed that there is a set extending this containment representation if and only if the original instance of 3SAT has a satisfying truth assignment. The idea is that any such set S must contain the element 0, and so S cannot be contained in $L \setminus c_i$ for any clause c_i , which is exactly the requirement that S contains a literal in c_i . In addition, the truth assignment given by S must form a valid partial truth assignment, since S can contain at most one of each pair of literals, as it cannot contain any set S_{v_i} . The other direction is again similar to the proof of Theorem 18, as the set of all literals a satisfying truth assignment makes true is a valid extension of the containment representation. \square

4.2 Containment-Free Representations

In the remainder of this section, we consider the problem of finding a minimum overlap representation where no set is contained in any other, or where the number of set containments is limited.

Problem. The CF-OVERLAP NUMBER problem is defined as:

Instance: A graph, $G = (V, E)$, and a natural number k .

Question: Does the graph G have a containment-free overlap representation of size k ?

In the absence of containment, the definitions of overlap and intersection coincide, and so this problem is equivalent to the problem of finding a minimum containment-free intersection representation. In order to show the hardness of this problem, we reduce the INTERSECTION NUMBER problem to it, since INTERSECTION NUMBER is known to be **NP**-complete [9].

Theorem 20. CF-OVERLAP NUMBER is **NP**-complete.

Proof. Given an instance $G = (V, E)$ and k of INTERSECTION NUMBER, with $n = |V|$, we construct the graph G' by adding, for each $v \in V$, a new vertex v' that is adjacent only to v . Let V' be the set of all new vertices in G' , and let E' be the set of all edges in G' incident on a vertex in V' . The instance of CF-OVERLAP NUMBER is then given by G' and $k + 2n$, which can clearly be constructed in polynomial time.

Notice that any containment-free overlap representation forms a containment free intersection representation, and further that in any containment-free intersection representation for G' , the sets S_v and $S_{v'}$ associated with a vertex $v \in V$ and $v' \in V'$ must share a common element, as these vertices are adjacent, and furthermore, since v' is adjacent only to v , this element is only be found in S_v and $S_{v'}$. The set $S_{v'}$ is not contained in S_v , and so it must contain at least one other element, which is unique to the set $S_{v'}$, since v' is adjacent only to v . This implies that for all $v \in V$, there are at least two elements found only in one or both of S_v and $S_{v'}$, which ensures that there are no containment relationships between any sets of the representation. Since these elements suffice to represent the vertices in V' , and the representation is already containment free, the remaining elements of the representation form an arbitrary intersection representation for G . Hence, the containment-free overlap number of G' is exactly $\theta_e(G) + 2n$, where $\theta_e(G)$ is the size of a minimum intersection representation for G . Thus G has an intersection representation of size k if and only if G' has a containment-free overlap representation of size $k + 2n$. \square

4.3 Overlap Representations with Limited Containment

We can extend the hardness of the CF-OVERLAP NUMBER problem to the problem of finding a minimum overlap representation of a graph, using at most a constant number of containment relationships between sets of the representation. Formalized as a decision problem, we consider the following problem. The factor of 2 appears since the nonadjacent pairs (u, v) and (v, u) are both counted, but we refer to the single non-edge as a containment relationship.

Problem. The L -CONTAINMENT OVERLAP NUMBER problem, for any natural number L , is defined as:

Instance: A graph, $G = (V, E)$, and a natural number k .

Question: Is there is some collection $\mathcal{C} = \{S_v : v \in V\}$ that forms an overlap representation, such that $|\bigcup_{v \in V} S_v| \leq k$ and $|\{(u, v) \notin E : u \neq v \text{ and } S_u \cap S_v \neq \emptyset\}| \leq 2L$?

This problem, when $L = 0$ is exactly the CF-OVERLAP NUMBER problem and so by Theorem 20 it is **NP**-complete in this case. For any constant L , a simple Turing reduction from the CF-OVERLAP NUMBER problem is given by making $2L+1$ copies of the input graph, and then finding an overlap representation with no more than L containments, which, by the pigeonhole principle, must leave at least one copy of G containment free, both internally, and with respect to other components of the graph. Furthermore, if we have a minimum representation, then this representation for G must also be minimum, as the sets associated with the vertices of this copy of G are disjoint from the sets associated with vertices in any other copy. With a little more work, we can find a many-one reduction from the CF-OVERLAP NUMBER problem, by adding to the graph G extra components where a minimum representation is be compelled to “spend” all L set containments, leaving G with a containment-free representation. We can do this in such a way that we can track the number of elements these extra components add to the representation. To show this result we make use of Corollary 6, which gives an upper bound on the number of elements we are able to save by allowing containment relationships between components of the constructed graph.

Theorem 21. *For any $L \in \mathbb{N}$, L -CONTAINMENT OVERLAP NUMBER is **NP**-complete.*

Proof. Let $G = (V, E)$ and k be an instance of CF-OVERLAP NUMBER. We set $n = |V|$, and we consider only cases where $n \geq 4$, as smaller cases can be solved as part of the transformation by searching all possible representations and producing as output a trivial yes or no instance. In the instance we construct, we add $2L$ components to the graph G . Each of these components is given by the graph $B_i = (V_i, E_i)$, which is constructed from $n + 1$ disjoint edges, with three nonadjacent universal vertices, as shown in Figure 2. More formally, the vertices V_i and the edges E_i of each component B_i are given by

$$\begin{aligned} V_i &= \{v_{i,j} : 1 \leq j \leq 2n + 2\} \cup \{x_i, y_i, z_i\}, \\ E_i &= \{(v_{i,2j-1}, v_{i,2j}) : 1 \leq j \leq n + 1\} \cup \\ &\quad \{(x_i, v_{i,j}), (y_i, v_{i,j}), (z_i, v_{i,j}) : 1 \leq j \leq 2n + 2\}. \end{aligned}$$

The graph in the constructed instance of L -CONTAINMENT OVERLAP NUMBER is then given by a disjoint union, $H = G + B_1 + B_2 + \dots + B_{2L}$, of $2L$ of these new components with the graph G . The value k' is set to

$$k' = k + 3L(n + 1) + 4L(n + 1), \tag{3}$$




Figure 2: Example of B_i with $n = 4$.

to complete the instance of L -CONTAINMENT OVERLAP NUMBER.

Before showing that the given instance of the CF-OVERLAP NUMBER problem is equivalent to the constructed instance, we first make some observations about overlap representations of the graphs B_i . In a minimum containment-free overlap representation for the vertices $v_{i,j}$ of B_i , we must use $3(n+1)$ elements, as each disjoint edge $(v_{i,2j-1}, v_{i,2j})$ requires at least three new elements in the representation. Furthermore, these three elements are given by an element unique to $S_{v_{i,2j-1}}$, an element unique to $S_{v_{i,2j}}$, and an element in the intersection of these two sets. We can extend a minimum representation for these vertices to include x_i and y_i without increasing the size of the representation. To do this, we set S_{x_i} to be those elements in common to the sets associated with both endpoints of each edge $(v_{i,2j-1}, v_{i,2j})$, and we set S_{y_i} to the elements that are unique to each of these sets. Since $n+1 \geq 2$, these sets are not be contained in any set $S_{v_{i,j}}$, and so these sets overlap, as desired. This forms the unique (up to permutation of the elements) minimum containment-free representation for all the vertices of B_i except z_i . If we allow a single containment relationship, we can set $S_{z_i} = S_{x_i}$ to obtain a representation with size $3(n+1)$.

If we seek a containment-free overlap representation for B_i the situation is more bleak, as we cannot extend the unique minimum containment representation for every vertex except z_i without adding new elements. This is because we still must use three elements to represent each edge $(v_{i,2j-1}, v_{i,2j})$, but there is no partition of these elements into three sets such that each set overlaps both $S_{v_{i,2j-1}}$ and $S_{v_{i,2j}}$. We are required then to use four elements for each edge $(v_{i,2j-1}, v_{i,2j})$, with two elements in common to the sets associated with the endpoints, which brings the size of a minimum containment-free overlap representation of B_i to $4(n+1)$. The key to the remainder of the proof is that by allowing a single containment relationship, we can reduce the size of the representation for some component B_i by $n+1$ elements.

If \mathcal{C} is a minimum L -overlap representation for H , of size no more than $k' = k + 3L(n+1) + 4L(n+1)$, we show that G has a containment-free overlap representation of size not more than k . We claim that, as \mathcal{C} is minimum, the representation \mathcal{C} when restricted to G is already containment-free, and in fact, the L containment relationships can be found in L of the components B_i . To show this, we examine the other potential cases for a non-edge to be represented by containment, showing in each one that we can make a local transformation to move the containment relationship to some component B_i , in the process reducing the size of the representation, contradicting the optimality of \mathcal{C} .

The first such case we consider is any containment within the representation of G , which is,

two vertices u and v such that $S_u \subseteq S_v$. We replace S_v with $n - 1$ new elements, a_1, a_2, \dots, a_{n-1} , to obtain $S'_v = \{a_1, a_2, \dots, a_{n-1}\}$. This removes the containment relationship between u and v , and forces S'_v not to contain any other set in the representation. In order to ensure that the representation is still valid, we modify the sets associated with some of the other vertices. There are exactly three ways a set can interact with S'_v : we can have the set we consider contain S'_v , the two sets can be disjoint, or the two sets can overlap. We consider, for each of these three interactions, how to alter the set to maintain a valid overlap representation of H . For any vertex w with $S_v \subseteq S_w$, we replace the set S_w with the set $S'_w = S_w \cup S'_v$, to ensure that this containment relationship is not altered. This alteration does not affect the overlap, containment, or disjointedness relationships of the set S_w , as these are new elements, and by transitivity, we have added these new elements to any set that contains S_w . If w is a vertex such that S_w and S_v are disjoint, then S_w and S'_v must also be disjoint, and there is nothing to do in this case. If w is such that S_w and S_v overlap, then we have $S'_v \cap S_w = \emptyset$, which we correct by setting $S'_w = S_w \cup \{a_i\}$, for an element $a_i \in S'_v$ that we have not already used for this purpose. This forces S'_w and S'_v to overlap, as the conditions that $n \geq 4$ and S_w overlaps the set S_v ensure that there are at least two elements in each of these sets. We must also add the element a_i to any set that contains S_w to preserve this containment relationship. This does not affect the representation of any vertex but v , as the sets that the element a_i is being added to must also intersect S_v , and we do not add all of the a_i to a set that should not contain S'_v , since there are at most $n - 2$ vertices that are adjacent to v . Thus, we can remove at least one containment relationship from G , by adding $n - 1$ new elements to the representation. Since there are $2L$ components B_i , and only $L - 1$ remaining containments, there must be some i for which the vertices of B_i are involved in no containment relationships. We can use the containment we just removed from G to reduce the size of the representation for B_i from $4(n + 1)$ to $3(n + 1)$, which, in total, saves at least $n + 1 - (n - 1) = 2$ elements from the representation, contradicting the assumption that \mathcal{C} was minimal. Thus, the vertices of G are not involved in any containment relationships in \mathcal{C} .

The second case of a containment relationship is one internal to one of the components B_i . If this containment is between two vertices $v_{i,j}$ and $v_{i,k}$, we can simply replace the representation for $v_{i,j}$ and the vertex $v_{i,j \pm 1}$ it forms an edge with. This is done by setting $S_{v_{i,j}} = \{a_1, a_3, a_4\}$ and $S_{v_{i,j \pm 1}} = \{a_2, a_3, a_4\}$, where the elements a_i are new to the representation. Finally, we add a_1 and a_2 to S_{x_i} , a_3 to S_{y_i} , and a_4 to S_{z_i} , being careful to add these elements to any set that contains these elements. If preserving these containment relationships results in all of $\{a_1, a_2, a_3, a_4\}$ being contained in one of S_{x_i}, S_{y_i} , or S_{z_i} , we simply add a_3 to each of S_{x_i}, S_{y_i} and S_{z_i} , and remove a_1, a_2 , and a_4 from these sets, once again being careful to preserve any containment relationships. This replacement removes the containment between $v_{i,j}$ and $v_{i,k}$, and leaves a valid overlap representation. As the cost of this alteration was only four elements, and we can apply the freed containment relationship to some other component B_j to save $n + 1$ containments, this also contradicts the optimality of \mathcal{C} .

The only remaining case for a containment internal to B_i is one between two of x_i, y_i , and z_i , as these vertices are adjacent to all other vertices of B_i . We must also consider the case that between the vertices x_i, y_i and z_i there are two or more containments, but since we can extend a minimum representation for the vertices $v_{i,j}$ to these vertices using only one containment, we can again apply this containment elsewhere, contradicting the optimality of \mathcal{C} .

The final case we must consider is a containment relationship between two vertices in differing components of H . Let U and W be the vertex sets of the two components, where for some vertex $u \in U$ and $w \in W$ we have $S_u \subseteq S_w$. Let $A = \bigcup_{u \in U} S_u$. Lemma 5 implies that for any vertex

in $v \in W$, either S_v contains A or it is disjoint from it. The elements of A then, within W , act as a single element. This allows these elements to be replaced with a single new element, where once again, whenever we add a new element to a set we must also add this new element to any sets that contained the original set. After this replacement has been made, we have removed at least one containment relationship, at a cost of one new element in the representation, which once again contradicts the optimality of \mathcal{C} .

Thus, a minimum L -containment overlap representation for H uses containment only between the vertices x_i, y_i , and z_i , and uses at most one containment per triple of vertices. Thus, in a minimum overlap representation, we have a containment-free overlap representation for G , and L of the B_i , and we have an overlap representation using only one containment for the remaining L of the B_i . Then, where r is the containment-free overlap number of G , this representation has size $r + 4L(n + 1) + 3L(n + 1)$, which by Equation (3) is less than k' only when $r \leq k$, as desired.

Fortunately, the other direction is simple. If we take any containment-free overlap representation for G of size no more than k , we can form the representations discussed above for each B_i , by simply using three elements per edge $(v_{i,2j-1}, v_{i,2j})$ for L of the B_i and four elements per edge for the remaining L . Placing the containments in appropriate places, we can find an L -containment overlap representation for H of size no more than $k + 3L(n + 1) + 4L(n + 1) = k'$, as required. \square

5 Conclusion

There are many open problems related to the overlap number of a graph. Foremost among these unanswered questions is the complexity of computing the overlap number.

Problem. The OVERLAP NUMBER problem is defined as:

Instance: A graph, $G = (V, E)$, and an integer k .

Question: Is there an overlap representation $\mathcal{C} = \{S_v : v \in V\}$ of G with $|\bigcup_{v \in V} S_v| \leq k$?

This problem is clearly in **NP**, as it is a simple matter to verify that a given representation is both correct and of the appropriate size, and the evidence suggests that this problem is also complete for **NP**. There are also many class of graphs for which no algorithm to find a minimum overlap representation is known. Many of these classes, such as cographs, are classes of graphs for which many other combinatorial problems are tractable, and so there is reason to believe that efficient algorithms exist to compute the overlap number on some such classes of graphs, but they have yet to be discovered.

Acknowledgements

This research was partially supported by NSERC and iCORE.

References

- [1] B. Bollobás. *Modern Graph Theory*. Springer, 1998.

- [2] B. Burlingham and L. Stewart. A translation of Sur deux propriétés des classes d'ensembles by Edward Szpilrajn-Marczewski. <http://webdocs.cs.ualberta.ca/~stewart/Pubs/MarczewskiTranslation.pdf>, 2009.
- [3] D. W. Cranston, N. Korula, T. D. LeSaulnier, K. G. Milans, C. Stocker, J. Vandenbusschek, and D. B. West. Overlap number of graphs. Submitted for publication, 2010.
- [4] P. Erdős, A. W. Goodman, and L. Pósa. The representation of a graph by set intersections. *Canadian Journal of Mathematics*, 18:106–112, 1966.
- [5] D. R. Fulkerson and O. A. Gross. Incidence matrices and interval graphs. *Pacific Journal of Mathematics*, 15(3):835–855, 1965.
- [6] M. C. Golumbic and E. R. Scheinerman. *Containment Graphs, Posets, and Related Classes of Graphs*, pages 192–204. Number 555 in Annals of the New York Academy of Sciences. New York Academy of Sciences, 1989.
- [7] P. Henderson. Overlap number lower bounds. Submitted for publication, 2010.
- [8] D. E. Knuth. *The Art of Computer Programming*, volume 1, Fundamental Algorithms. Addison-Wesley, third edition, 1997.
- [9] L. T. Kou, L. J. Stockmeyer, and C. K. Wong. Covering edges by cliques with regard to keyword conflicts and intersection graphs. *Communications of the ACM*, 21(2):135–139, 1978.
- [10] E. C. Milner. A combinatorial theorem on systems of finite sets. *Journal of the London Mathematical Society*, 43:204–206, 1966.
- [11] E. Prisner. Graphs with few cliques. In *Proceedings of the 7th Quadrennial International Conference on the Theory and Applications of Graphs, 1992*, Graph Theory, Combinatorics, and Applications, pages 945–956, 1995.
- [12] A. Raychaudhuri. Intersection number and edge clique graphs of chordal and strongly chordal graphs. *Congressus Numeratium*, 67:197–204, 1988.
- [13] A. Raychaudhuri. Edge clique graphs of some important classes of graphs. *Ars Combinatorica*, 32:269–278, 1991.
- [14] W. Rosgen. Set representations of graphs. Master's thesis, University of Alberta, 2005.
- [15] T. J. Schaefer. The complexity of satisfiability problems. In *Proceedings of the 10th Annual ACM Symposium on Theory of Computing*, pages 216–226, 1978.
- [16] E. Szpilrajn-Marczewski. Sur deux propriétés des classes d'ensembles. *Fundamenta Mathematicae*, 33:303–307, 1945.

Appendix

This appendix contains the proofs that have been omitted from the main text.

Proofs Omitted From Section 3

Lemma 11. For $1 \leq k < n$,

$$\binom{n}{k} \geq \sqrt{\frac{1}{8\pi k}} \left(\frac{n}{k}\right)^k \left(\frac{n}{n-k}\right)^{n-k}.$$

Proof. By simple expansion, using Stirling's Approximation (Equation (1)), we have

$$\begin{aligned} \binom{n}{k} &= \frac{n!}{k!(n-k)!} \geq \frac{\sqrt{2\pi n}(n/e)^n}{2\pi e^{1/(12k)+1/(12(n-k))} \sqrt{k(n-k)}(k/e)^k((n-k)/e)^{n-k}} \\ &\geq \frac{1}{e^{1/6}} \sqrt{\frac{n}{2\pi k(n-k)}} \frac{n^n}{k^k(n-k)(n-k)} \\ &\geq \frac{1}{2} \sqrt{\frac{n}{2\pi k(n-k)}} \left(\frac{n}{k}\right)^k \left(\frac{n}{n-k}\right)^{n-k} \\ &\geq \sqrt{\frac{n-k}{8\pi k(n-k)}} \left(\frac{n}{k}\right)^k \left(\frac{n}{n-k}\right)^{n-k} \\ &= \sqrt{\frac{1}{8\pi k}} \left(\frac{n}{k}\right)^k \left(\frac{n}{n-k}\right)^{n-k} \end{aligned}$$

as in the statement of the lemma. □

Theorem 12. For $n \geq 1$,

$$\min \left\{ m : n \leq \binom{m}{\lfloor \frac{m+2}{2} \rfloor} \right\} \in \Theta(\log n).$$

Proof. Let $x = \min \left\{ m : n \leq \binom{m}{\lfloor \frac{m+2}{2} \rfloor} \right\}$. We show a lower bound by observing that there are 2^x subsets of $\{1, 2, \dots, x\}$, and so we must have $n < 2^x$, which implies that $x \in \Omega(\log n)$. Turning to an upper bound, notice that, by the definition of x , we have $\binom{x-1}{\lfloor \frac{x+1}{2} \rfloor} < n$. Using this, and Lemma 11, we have

$$\begin{aligned} n &> \binom{x-1}{\lfloor \frac{x+1}{2} \rfloor} \geq \binom{x-1}{\frac{x+1}{2}} \geq \sqrt{\frac{2}{8\pi(x+1)}} \left(\frac{2(x-1)}{x+1}\right)^{(x+1)/2} \left(\frac{2(x-1)}{x-1}\right)^{(x-1)/2} \\ &= 2^x \sqrt{\frac{1}{4\pi(x+1)}} \left(\frac{x-1}{x+1}\right)^{(x+1)/2} \geq 2^{(x-1)/2} \sqrt{\frac{1}{4\pi(x+1)}} \end{aligned}$$

We can then take logarithms to obtain

$$\begin{aligned} \log n &> \log \left(2^{(x-1)/2} \sqrt{\frac{1}{4\pi(x+1)}} \right) > \frac{x-1}{2} + \frac{1}{2} \log \left(\frac{1}{4\pi(x+1)} \right) \\ &= \frac{x-1}{2} - \frac{\log(4\pi(x+1))}{2} = \frac{x-1}{2} - \frac{\log 4\pi + \log(x+1)}{2} \end{aligned}$$

Since $x \in \Omega(\log n)$, for large enough n we must have $\log(x+1) < x/2$. We then have, by the above, and setting $C = \log(4\pi)/2$,

$$\log n > \frac{x-1}{2} - \frac{x}{4} - C = \frac{x-2}{4} - C,$$

which is $x/4 < \log n + 1/2 + C$, and so we have $x \in O(\log n)$, as desired. □